

Review Article

Role of BDNF-TrkB Signaling in Regulating Anxiety and Depression-Like Behavior in Diverse Brain Regions

Yafei Shi, Chunguang Mu, Yongjuan Xin*

School of Public Health, Zhengzhou University, Zhengzhou, China

Email address:

syf20201010@163.com (Yafei Shi), 15738221749@163.com (Chunguang Mu), yjxinzzu@163.com (Yongjuan Xin)

*Corresponding author

To cite this article:Yafei Shi, Chunguang Mu, Yongjuan Xin. Role of BDNF-TrkB Signaling in Regulating Anxiety and Depression-Like Behavior in Diverse Brain Regions. *Clinical Neurology and Neuroscience*. Vol. 7, No. 2, 2023, pp. 18-30. doi: 10.11648/j.cnn.20230702.11**Received:** June 12, 2023; **Accepted:** June 27, 2023; **Published:** July 6, 2023

Abstract: Depressive disorders occur often jointly with anxiety disorders, which can cause serious health problems. The underlying mechanism is not fully understood. Figuring out the mechanism of depressive and anxiety disorders would benefit patients in future therapy. Brain-derived neurotrophic factor (BDNF) is a famous neurotrophin that modulates synaptic plasticity in the brain. It is generally believed that decreased BDNF levels are associated with depression. The purpose of this review is to elucidate the role of the BDNF-TrkB signaling pathway in different brain regions and its antidepressant effect, to provide scientific evidence for the treatment of anxiety and depression. The changes of the BDNF-TrkB signaling pathway before and after antidepressant treatment were compared by retrieving preclinical studies related to the BDNF-TrkB signaling pathway and classifying them according to different brain regions. It is found that the concentration of BDNF varies in different brain regions. The inhibition of the BDNF-TrkB pathway in the cortex, hippocampus, and amygdala and the activation of the BDNF-TrkB pathway in the anterior cingulate cortex (ACC), nucleus accumbens (NAc), and lateral habenula (LHb) is associated with anxiety and depression-like behaviors. Lacking BDNF or its receptor TrkB is not the cause of anxiety or depression, but affects the effect of antidepressant treatment. Increased BDNF can alleviate anxiety and depression. There are still other molecules that can regulate anxiety and depression-like behaviors by influencing the expression of BDNF or TrkB. The function of BDNF in the ACC, NAc, and LHb areas needs to be further explored.

Keywords: Depression, Anxiety, BDNF, TrkB, Mechanism, Pathway

1. Introduction

Depressive disorders are mental disorders characterized by the presence of a sad, empty, or irritable mood, accompanied by somatic and cognitive changes that significantly affect the individual's capacity to function [1]. Depressive disorders occur often jointly with anxiety disorders [2]. Around 85% of patients who have depression will also exhibit symptoms of anxiety, while up to 90% of individuals with anxiety disorders will also experience comorbid depression [3]. Depression and anxiety were two of the top ten causes of global disability-adjusted life-years (DALYs) in adolescents aged 10–24 years in 2019 [4]. About 4.7% of the world's population has an episode of depression every year [5]. The causes of depression and anxiety disorders are complex, including

genetic factors and environmental factors. People with high depression polygenic risk scores [6] are more susceptible to depressive disorders. Adverse experiences such as social isolation [7], trauma [6], and childhood maltreatment [8] were significantly associated with depressive disorders. During the pandemic of COVID-19, the global prevalence of major depressive disorders (MDD) and anxiety disorders increased [9]. Figuring out the underlying mechanism is important to treat these mental disorders effectively. The pathogenesis of depressive disorders is not fully understood. There are some hypotheses about depression, including the neurotrophic hypothesis, monoamine hypothesis, hypothalamic-pituitary-adrenal axis hypothesis, etc. [10].

Brain-derived neurotrophic factor (BDNF), which is widely expressed in the brain, is the most studied neurotrophic factor

and plays a key role in the central nervous system by supporting neuron survival and facilitating neurogenesis [11]. It performs biological functions by binding to its specific receptors. There are at least two BDNF-specific receptors, TrkB and P75^{NTR}, on the nerve cell membrane, and BDNF has a higher affinity with TrkB. Cavaleri D et al. found that people with MDD have lower peripheral and central BDNF levels than non-depressive individuals, based on meta-analyses [12].

To explore the molecular mechanisms of depression and anxiety and the effect of antidepressant treatment, experimental paradigms such as chronic unpredictable mild stress (CUMS), chronic restraint stress (CRS), learned helplessness (LH), corticosterone (CORT), lipopolysaccharide (LPS), etc. were used to induce animals showing anxiety and depression-like behaviors. In this review, we will elucidate the function of the BDNF-TrkB signaling pathway in diverse brain regions by giving a summary of the preclinical anxiety and depression-like behaviors studies.

2. BDNF-TrkB Signaling Pathway in Anxiety and Depression-Like Behavior

Impaired BDNF-TrkB signaling is associated with anxiety and depression-like behaviors (Figure 1). BDNF levels in MDD patients changed before and after antidepressant treatment. The studies on MDD patients’ post-mortem brain samples found that the levels of BDNF were decreased in the hippocampus [13] and amygdala [14]. After the effective treatment of antidepressant drugs [15] or electroconvulsive therapy (ECT) [16], the serum BDNF levels increased. And it was also found that BDNF levels in the hippocampus increased in antidepressant-treated MDD patients’ brain samples after death [17]. It suggested that decreased BDNF concentration was associated with anxiety and depression-like behavior, and preclinical studies also proved this and found something new. The relation between BDNF-TrkB signaling and anxiety and depression-like behavior in different brain regions was not consistent. (Table 1)




Figure 1. Impaired BDNF-TrkB signaling pathway in anxiety and depression-like behaviors.

Table 1. The alteration of BDNF and TrkB in different brain regions in anxiety and depression-like behavior.

| Pathway | Animal | Paradigm | Phenotype | | Brain regions | Before treatment | | After treatment | |
|-----------------------|--------|----------|------------------------|---------------------------|---------------|------------------|------|-----------------|------|
| | | | Anxiety-like behaviors | Depression-like behaviors | | BDNF | TrkB | BDNF | TrkB |
| BDNF | Rats | LPS | - | - | CT, Hip | ↓ | - | - | - |
| BDNF | Mice | CUMS | Yes | Yes | CT, Hip | ↓ | - | - | - |
| BDNF-TrkB | Rats | CRS | - | Yes | CT, Hip | ↓ | ↓ | ↑ | ↑ |
| PIK3CA-AKT1-NRF2/BDNF | Rats | OBX | Yes | Yes | CT, Hip | ↓ | ↓ | ↑ | ↑ |
| Nrf2/BDNF | Mice | CORT | Yes | Yes | CT, Hip | ↓ | - | ↑ | - |
| BDNF | Rats | LH | - | - | PFC, Hip | ↓ | - | - | - |

| Pathway | Animal | Paradigm | Phenotype | | Brain regions | Before treatment | | After treatment | | |
|-----------------------------|----------|---|------------------------|---------------------------|-------------------|------------------|------|-----------------|------|------|
| | | | Anxiety-like behaviors | Depression-like behaviors | | BDNF | TrkB | BDNF | TrkB | |
| BDNF | Mice | LPS | Yes | Yes | PFC, Hip | - | - | ↑ | ↑ | [24] |
| BDNF-TrkB | Mice | LPS | - | Yes | PFC, Hip | ↓ | ↓ | - | ↑ | [25] |
| BDNF-TrkB | Rats | CUS | Yes | - | PFC, Hip | ↓ | ↓ | ↑ | ↑ | [26] |
| BDNF-TrkB | Rats | Immune system activation | Yes | Yes | PFC, Hip | ↓ | ↓ | - | - | [27] |
| BDNF-ERK-CREB | Mice | Methamphetamine withdraw | Yes | Yes | PFC, Hip | ↓ | - | - | - | [28] |
| BDNF/VEGF | Mice | Burn | Yes | Yes | PFC, Hip | ↓ | - | ↑ | - | [29] |
| BDNF-CREB | Rats | CUMS | Yes | Yes | PFC | ↓ | - | ↑ | - | [30] |
| BDNF-HCN1 | Rats | SPS&S | Yes | Yes | PFC | ↓ | - | ↑ | - | [31] |
| mTOR-BDNF | Rats | CUMS | Yes | Yes | PFC | ↓ | - | ↑ | - | [32] |
| BDNF-TrkB-Akt | Mice | Subchronic arsenic exposure combined with reserpine | Yes | Yes | PFC | ↓ | ↓ | - | - | [33] |
| TrkB/Hsp70 | Hamsters | CUMS | Yes | Yes | PFC, Hip, Amy, Ht | - | ↓ | - | ↑ | [34] |
| BDNF | Rats | LPS/IL-1 β | - | - | Hip | ↓ | - | - | - | [35] |
| BDNF | Rats | CUMS | Yes | Yes | Hip | ↓ | - | - | - | [36] |
| BDNF | Rats | CUMS | - | Yes | Hip | ↓ | - | ↑ | - | [37] |
| BDNF | Rats | EM | Yes | Yes | Hip | ↓ | - | - | - | [38] |
| BDNF | Mice | HFD | No | Yes | Hip | ↓ | - | - | - | [39] |
| BDNF | Mice | FST | - | Yes | Hip | - | - | ↑ | - | [40] |
| BDNF | Mice | CD | Yes | Yes | Hip | ↓ | - | - | - | [41] |
| BDNF-TrkB | Mice | A β ₁₋₄₂ | Yes | Yes | Hip | ↓ | ↓ | ↑ | ↑ | [42] |
| BDNF-CREB | Rats | PM2.5 | Yes | Yes | Hip | ↓ | - | - | - | [43] |
| BDNF-TrkB-CREB | Mice | CORT | No | Yes | Hip | ↓ | ↓ | ↑ | ↑ | [44] |
| BDNF-TrkB-CREB | Mice | Noise | Yes | Yes | Hip | ↓ | ↓ | ↑ | ↑ | [45] |
| BDNF-TrkB-CREB | Mice | CRS | Yes | Yes | Hip | ↓ | - | ↑ | ↑ | [46] |
| BDNF-TrkB-CREB | Rats | CUS | Yes | Yes | Hip | ↓ | - | ↑ | - | [47] |
| BDNF-TrkB-CREB | Rats | CUS | Yes | Yes | Hip | ↓ | ↓ | - | - | [48] |
| BDNF-TrkB-CREB-E RK | Mice | CUMS | Yes | Yes | Hip | ↓ | ↓ | ↑ | ↑ | [49] |
| PGC-1 α -FNDC5-BD NF | Rats | CUMS | Yes | Yes | Hip | ↓ | - | ↑ | - | [50] |
| mTOR-BDNF | Mice | A β ₁₋₄₀ | Yes | Yes | Hip | ↓ | - | ↑ | - | [51] |
| AMPK/BDNF | Mice | IBD | - | Yes | Hip | ↓ | - | ↑ | - | [52] |
| BDNF-TrkB-Akt-Gsk 3b | Rats | CUMS | Yes | - | Ht | - | ↓ | - | ↑ | [53] |
| CREB-BDNF | Mice | LPS | Yes | Yes | ACC | ↑ | - | ↓ | - | [54] |
| NF- κ B/BDNF | Mice | Hypoxia | Yes | Yes | Amy | ↓ | - | ↑ | - | [55] |
| BDNF-TrkB-CRH | Rats | Alcohol | Yes | - | Amy | ↓ | ↓ | ↑ | ↑ | [56] |
| BDNF | Rats | LH | - | - | NAC | ↑ | - | - | - | [23] |
| BDNF | Mice | CSDS | - | - | NAC | ↑ | - | ↓ | - | [57] |
| BDNF-TrkB | Mice | LPS | - | Yes | NAC | ↑ | ↑ | - | ↓ | [25] |
| BDNF | Rats | LH/Fst | - | Yes | DRN | - | - | ↑ | - | [58] |
| BDNF | Mice | CRS | Yes | Yes | LHb | ↑ | - | ↓ | - | [59] |

Abbreviations: ACC, Anterior cingulate cortex; Amy, Amygdala; CD, Crohn's disease; CORT, Corticosterone; CRS, Chronic restraint stress; CSDS, Chronic social defeat stress; CT, Cortex; CUMS, Chronic unpredictable mild stress; CUS, Chronic unpredictable stress; DRN, Dorsal raphe nucleus; EM, Endometriosis model; FST, Forced swimming test; HFD, High-fat diet; Hip, Hippocampus; Ht, Hypothalamus; IBD, Inflammatory bowel disease; LH, Learned helplessness; LHb, Lateral habenula; LPS, Lipopolysaccharide; NAC, Nucleus accumbens; OBX, Olfactory bulbectomy; PFC, Prefrontal cortex; SPS&S, Single prolonged stress & electric foot shock.

2.1. Cortex

The cortex is the layer of gray matter on the surface of the brain. It occupies about 80% of the brain and is responsible for perception, thinking, language, memory, etc. BDNF can promote neuron growth and the formation of synapses in the brain. The inhibition of the BDNF-TrkB signaling in the cortex is correlated with anxiety and depression. Anxiety and depression-like behavior [19, 22, 24, 32] can be induced by CUMS, LPS, CORT, etc. The BDNF [18, 25, 30, 32] and TrkB [26, 34] levels were decreased in the cortex. With BDNF decreased, dendritic spine density was also decreased in the cortex, and rats were susceptible to depression-like behavior in the LH [23].

2.2. Hippocampus

The hippocampus is beneath the cerebral cortex, and in the medial temporal lobe of the brain. It occupies a little volume of the brain and is responsible for learning, memory, and emotion. The dentate gyrus (DG), CA1, CA2, and CA3 regions of the hippocampus were investigated in many studies. The inhibition of the BDNF-TrkB signaling in the hippocampus is associated with anxiety and depression. In addition to the above-mentioned CUMS, CORT, LPS-induced emotional disorder models, chronic restraint stress (CRS), olfactory bulbectomy (OBX), A β ₁₋₄₀, A β ₁₋₄₂-induced model, and DSS-induced IBD model, can also lead rodents exhibiting

anxiety and depression-like behaviors [21, 42, 46, 51, 52]. The expressions of BDNF and TrkB [19, 21, 44, 47] in the hippocampus were decreased. And to be specific, BDNF mRNA levels in the dorsal hippocampus were decreased in the CRS model [46]. BDNF levels and dendrite spine density in CA3 and DG regions of the hippocampus of the LH susceptible group were significantly decreased [23]. Inflammation can lead to anxiety and depression-like behaviors. LPS model or DSS-induced IBD model can cause systemic inflammation including brain inflammation. BDNF was decreased and the inflammatory factors elevated [18, 25] in the hippocampus, especially in CA2 [35], CA3 [25, 35], and DG [25, 35] in LPS-induced models. Decreased proliferating cells and activation of astrocytes and microglia could be seen in the IBD model [52]. The proliferating and survival cells decreased in the ventral, dorsal, and DG of the hippocampus [51], and autophagy cells increased [42] in the A β -induced models. CUMS cause decreasing BDNF and TrkB and has an influence on synaptic plasticity, which shows by the decreases of PSD95 level and synaptic spines in the hippocampal CA1 region, and the apoptosis of GFAP positive cells increased [47]. Early postpartum exposure to high doses of PM2.5 induced anxiety and depression-like behaviors in rats, impairing the synaptic number in the hippocampal CA1 region, and the synaptic plasticity in the hippocampal region with PSD95 and SYP decreased [43].

2.3. Hypothalamus

The hypothalamus is located under the dorsal thalamus and accounts for 0.3% weight of the brain. It involves the regulation of body temperature, feeding, endocrine, etc. The inhibition of the BDNF-TrkB signaling in the hypothalamus is associated with anxiety and depression. In CUMS paradigm, decreased TrkB mRNA in the VMH area of the hypothalamus was correlated with anxiety and depression-like behaviors [34]. Anxiety-like behavior can be seen in the acute sleep deprivation (SD) model, with decreased TrkB and p-Akt levels, elevated serum corticosterone and inflammation factor, and astrocyte activation in the hypothalamus [53].

2.4. Amygdala

The amygdala is located at the anterior end of the inferior horn of the lateral ventricle. It is mainly involved in memory, visceral and endocrine regulation, emotional activity, etc. The inhibition of the BDNF-TrkB signaling in the amygdala is associated with anxiety and depression. Decreased BDNF and p-TrkB in the amygdala were observed in the hypoxic stress and repeated alcohol exposure-induced models [55, 56]. In the hypoxic stress model, the brain exhibits oxidative stress and inflammation status, with increased MDA, decreased antioxidants like GSH, SOD, and CAT, and increased inflammatory factors [55]. And the serum corticosterone was elevated with the increase of CRH-positive cells in the PVN of the hypothalamus [56].

2.5. Anterior Cingulate Cortex (ACC)

ACC is located in the parahippocampal gyrus region and

regulates emotional function. Activation of the ACC region and elevated BDNF levels are associated with anxiety and depression. In LPS-induced acute stress in rats, the ACC region was activated with an increasing level of c-Fos, and CREB, p-CREB, and BDNF levels were elevated in ACC [54]. Anxiety and depression-like behaviors can also occur in spared nerve injury (SNI) model. In this model, the ACC region was activated, and the expressions of BDNF and CREB were increased [54]. By knocking out CREB in ACC specifically, the expressions of CREB, p-CREB, and BDNF were all reduced, while the expressions of these proteins in the spinal cord were all increased, which could reduce anxiety and depression in rats [54]. With the silence of the ACC region, anxiety, and depression-like behaviors could be alleviated in rats [54]. The anti-BDNF antibody injection in bilateral ACC did not reverse the depression-like behavior of CUMS-induced mice [19]. This indicated that the expression of BDNF can be regulated by CREB.

2.6. Ventral Tegmental Area - Nucleus Accumbens (VTA-NAc)

VTA is located in the medial substantia nigra of the midbrain and the dorsal interpeduncular nucleus. NAc is located at the junction of basal ganglia and limbic system. VTA-NAc projecting has been proven in many studies [60, 61]. The activation of the BDNF-TrkB signaling in the VTA-NAc is associated with anxiety and depression. In the CSDS paradigm, the c-Fos protein was elevated in the VTA-NAc, and BDNF was increased in NAc [57]. Increased BDNF levels in the NAc can also be seen in LPS or LH-induced models [23, 25]. Meanwhile, the density of dendritic spines was also found higher than the control group, which indicated that regional differences in BDNF levels and dendritic spine density in the brain may be helpful to resist inevitable stress [23].

2.7. Dorsal Raphe Nucleus (DRN)

DRN is located in the midbrain and is the most important nucleus of serotonin neurons in the central nervous system. The inhibition of the BDNF-TrkB signaling in the DRN is associated with anxiety and depression. Infusion of BDNF in the midbrain near the DRN could reverse the depression-like behavior of rats [58]. Conditional knockout BDNF or TrkB in the DRN could not lead to anxiety or depression-like behavior, but conditional knockout TrkB could lead to loss of antidepressant efficacy, not BDNF [62]. Thus, we can see that TrkB in DRN may play an important role during antidepressants.

2.8. The Lateral Habenula (LHb)

The habenula is located above the posterior thalamus near the midline. It consists of the LHb and the medial habenula (MHb). The activation of the BDNF-TrkB signaling in the LHb is associated with anxiety and depression. In the CRS-induced model, the LHb was activated with an increased expression of c-Fos, and BDNF [59]. One study showed that

effort-based reward (EBR) contingency training could be a behavioral therapeutic intervention for depressive symptoms in rats, and the expression of BDNF in LHb decreased in EBR contingency training rats [63, 64]. After the BDNF was knockout in the LHb, CRS could not induce anxiety and depression-like behaviors in mice, and the expression of c-Fos and BDNF were not increased in the CRS paradigm [59].

3. The Function of BDNF in Antidepressant Treatment

In the combination of CUMS and 12-hour sleep deprivation (SD)-induced model, the depression-like behavior was alleviated with the hippocampal BDNF increased, but the anxiety-like behavior did not improve [49]. Conditional knockout BDNF in CA1 or DG of the hippocampus and DRN of the midbrain could not induce anxiety or depression-like

behavior but could impair the antidepressant function [62, 65]. After the treatment of antidepressants, the BDNF cKO mice in CA1 or DG showed depression-like behavior compared with wild-type mice, but without anxiety-like behavior [65]. Conditional knockout TrkB in DRN also didn't show anxiety or depression-like behavior, and after the treatment of antidepressants, mice exhibited depression-like behavior but not anxiety-like behavior [62]. Injecting BDNF into DG, CA3 of the hippocampus, DRN of the midbrain, and VTA showed an antidepressant role but not an antianxiety function [58, 66, 67]. The injection of TrkB.T1 into NAc could lead to depression-like behavior, not anxiety [67]. Conditional knockout BDNF in LHb could prevent depression-like behavior but not anxiety-like behavior [59]. Thus, it seems like BDNF-TrkB signaling works weak in anxiety-like behaviors. (Table 2)

Table 2. Preclinical studies of BDNF-TrkB signaling in anxiety and depression-like behavior.

| Pathway | Animal | Paradigm | Before treatment | | After treatment | | |
|--------------------|--------|--------------------------|------------------------|---------------------------|------------------------|--|------|
| | | | Anxiety-like behaviors | Depression-like behaviors | Anxiety-like behaviors | Depression-like behaviors | |
| BDNF-TrkB-CREB-ERK | Mice | SD 12h after CUMS | Yes | Yes | Not improved | Improved | [49] |
| BDNF | Mice | BDNF cKO in DG | No | No | Not improved | Improved in wildtype mice. Not improved in BDNF cKO mice | [65] |
| BDNF-TrkB | Mice | BDNF cKO in DRN | No | No | Not improved | Improved | [62] |
| BDNF-TrkB | Mice | TrkB cKO in DRN | No | No | Not improved | Improved in wildtype mice. Not improved in TrkB cKO mice | [62] |
| BDNF | Rats | BDNF infusion in DG | No | Yes | Not improved | Improved | [66] |
| BDNF | Rats | BDNF infusion in CA3 | No | Yes | Not improved | Improved | [66] |
| BDNF | Rats | BDNF infusion in CA1 | No | Yes | Not improved | Not improved | [66] |
| BDNF | Rats | BDNF infusion in DRN | No | Yes | Not improved | Improved | [58] |
| BDNF-TrkB | Rats | BDNF infusion in NAc | - | - | No | Yes | [67] |
| BDNF-TrkB | Rats | TrkB. FL infusion in NAc | - | - | No | No | [67] |
| BDNF-TrkB | Rats | TrkB. T1 infusion in NAc | - | - | No | Improved | [67] |
| BDNF | Mice | BDNF cKO in LHb | Yes | Yes | Not improved | Improved | [59] |

Abbreviations: cKO, Conditional knockout; CUMS, Chronic unpredictable mild stress; DG, Dentate gyrus; DRN, Dorsal raphe nucleus; LHb, Lateral habenula; NAc, Nucleus accumbens; SD, Sleep deprivation.

Patients with endometriosis have a high incidence of anxiety and depression. By establishing the rat endometriosis model (EM), researchers found that MDA in the hippocampus of rats began to rise 14 days after EM, at which point depression-like behaviors began to appear. Until 21 days after EM, the rats' anxiety-like behaviors gradually increased, at which time corticosterone increased. The hippocampus had pro-oxidation status with increased TPO, decreased GSH and SOD activity, and low BDNF levels [38]. This indicates that in the EM model, decreased BDNF levels in the hippocampus lag behind anxiety and depression-like behaviors. And BDNF may be not the cause of anxiety and depression, it might be the mediator towards the progression or prevention of anxiety and depression.

This phenotype was also confirmed by the transgenic mice model. Most BDNF knockout mice died a few days after birth due to non-brain tissue effects such as cardiac dysplasia, but some individuals can survive for 2-4 weeks [68]. BDNF^{+/-}

mice didn't exhibit depression-like behavior [69]. No anxiety and depression-like behaviors were observed in BDNF conditional knockout mice in the forebrain, hippocampal CA1 and DG regions, and DRN of the midbrain, suggesting that the absence of BDNF in the hippocampus and the midbrain does not induce anxiety or depression-like behaviors [62, 65, 70, 71]. But the absence of BDNF in the forebrain and DG region of the hippocampus rather than the CA1 region plays a role in antidepressant treatment [65, 71], suggesting that genetic damage to BDNF-TrkB signaling pathways does not appear to lead to depression-like behavior, but will hinder the role of antidepressants [72]. Thus, BDNF may be a target for antidepressants, but not the only mediator of depression or anxiety.

3.1. Cortex

BDNF-TrkB is a target for antidepressant therapy. Duman's group first proposed that the common target of antidepressant

therapy may be BDNF and its receptor TrkB [20]. Researchers used electroconvulsive therapy, a variety of antidepressants, and a variety of non-antidepressant psychotropic therapies to treat the depression-like behavior of rats induced by CRS, to protect or reduce the damage of neurons by increasing the expression levels of BDNF and TrkB [20]. The following preclinical experiments also confirmed this. After treatment of anxiety and depression-like behavior induced by CUMS and LPS, the BDNF-TrkB signaling was activated, and the levels of BDNF and TrkB were increased [24-26], and anxiety and depression-like behaviors of rodents were improved. This suggests that the BDNF-TrkB pathway plays an important role in the pathophysiological processes of anxiety and depression-like behaviors. Antidepressant therapy was ineffective in BDNF^{+/-} transgenic mice also proved it [69]. During antidepressant therapy, BDNF mediated the activation of TrkB and induced TrkB autophosphorylation, which led to the phosphorylation of CREB [69]. In LPS induced acute stress model, intraperitoneal (i. p.) injection of TrkB agonist 7, 8-dihydroxyflavonoids (7, 8-DHF) showed antidepressant effects, significantly alleviating the reduction of p-TrkB in the prefrontal cortex, while pre-injection of TrkB antagonist ANA-12 blocked the antidepressant effects of 7, 8-DHF [25]. Compared with wild-type mice, cortical TrkB-T1 subtype overexpressing mice showed decreased TrkB signaling in the brain [73] and decreased responsiveness to antidepressant drugs [69]. These studies suggest that the BDNF-TrkB signaling pathway plays a role in depression-like behavior and antidepressant effects.

3.2. Hippocampus

The BDNF-TrkB pathway in the hippocampus plays an important role in mediating the effect of antidepressant therapy. BDNF and TrkB mRNA levels increased in the granule cell layer of the DG region and conical cell layer of the CA3 and CA1 region after antidepressant or electroconvulsive therapy but remained unchanged after non-antidepressant psychotropic medication [20]. Multiple studies proved that BDNF and TrkB increased after the improvement of anxiety and depression-like behaviors after antidepressant therapy [21, 44, 46]. The combination of transcranial photobiomodulation (tPBM) and housing in an enriched environment (EE) can decrease the corticosterone concentration and activate the BDNF-TrkB-CREB signaling pathway in a noise stress-induced model [45]. Continuous or interval exercise preconditioning could mitigate the anxiety and depression-like behaviors of CUMS by activating the PGC-1 α /FNDC5/BDNF pathway in the hippocampus [50]. Exercise can increase the number of hippocampal cell proliferation and survival in the A β -induced model [51]. Anti-inflammation treatment can activate the antioxidant factor Nrf2 signaling, to reduce inflammatory factors, and activate BDNF-TrkB signaling to improve anxiety and depression in LPS [24] and OBX-induced models [21]. In the CORT-induced model, fluoxetine treatment can activate the Nrf2 signaling pathway, and increase the BDNF expression to play an anti-anxiety depression effect [22]. But Nrf2 signaling was not the upstream signaling pathway of BDNF-TrkB signaling, for

Nrf2 knockout mice can present elevated BDNF expression [22]. Liver hydrolysate can prevent depressive-like behavior in the DSS-induced IBD model by activating the AMPK-BDNF signaling to increase the proliferation of hippocampal cells and neurogenesis [52]. These results suggest that BDNF and its receptor TrkB may be the target of antidepressant therapy.

To verify the antianxiety and antidepressant functions of BDNF-TrkB, researchers infused BDNF into the DG, CA3, and CA1 areas of the bilateral hippocampus respectively in the LH paradigm, then they found that the DG and CA3 areas were activated with the improvement of depression-like behavior, which was confirmed by the increment of c-Fos and Elk-1 [66]. Ketamine can alleviate anxiety and depression-like behaviors, and TrkB antagonist (K252a) can antagonize its therapeutic effect [47]. TrkB agonist 7, 8-dihydroxyflavone (7, 8-DHF) can exhibit antidepressant function by preventing the decrease of p-TrkB in DG and CA3 areas of the hippocampus in the LPS-induced model [25]. In the LH paradigm, inhibition of Trk (K252a) or MEK (U0126) blocked the antidepressant effect of BDNF, suggesting that the effect of BDNF is mediated by the activation of MEK and MAPK cascades [66].

3.3. Hypothalamus

After anti-inflammation treatment, with the PI3K-Akt and BDNF-TrkB signaling pathway in the hypothalamus activated, the neuroinflammation was alleviated, corticosterone concentration decreased, the activation of astrocytes in the hypothalamus was inhibited, and anxiety-like behavior was improved [53]. Infusing BDNF into the DG region of the hippocampus, TrkB mRNA in the VMH region of the hypothalamus was up-regulated, showing an antidepressant effect [34].

3.4. Amygdala

After the treatment of drugs or acupuncture, BDNF and p-TrkB were increased, and the anxiety and depression-like behaviors were alleviated [55, 56]. At the same time, the lipid peroxidation and inflammation levels in the brain and the serum corticosterone were decreased, with NF- κ B signaling inhibited [55]. The expression of CRH in the PVN also decreased [56]. Injecting BDNF into the DG area, TrkB mRNA increased in the CeA of the amygdala, which performed an antidepressant function [34]. In the LPS paradigm, IL-33 was up-regulated in the amygdala, and IL-33 knockout mice exhibited an antianxiety effect with BDNF level increased in the amygdala [74]. Injecting TrkB antagonists, ANA-12, in the amygdala can prevent the increase of CRH in PVN, and alleviate the anxiety-like behavior of rats [56]. This indicated that activating BDNF-TrkB signaling could decrease the corticosterone level by regulating the CRH-positive neurons in PVN.

3.5. VTA-NAc

BDNF protein in NAc was thought to be derived from VTA [75]. Infusing BDNF in VTA for 1 week, the latency of immobility was shortened by 57%, which means that the rats

showed depression-like behavior [67]. Bilateral NAc infusion of TrkB antagonist, ANA-12, showed an antidepressant effect by preventing the elevated p-TrkB level [25]. To study the function of TrkB in NAc, rats received an intra-NAc full-length (TrkB. FL) or truncated (TrkB. T1) TrkB AAV injection. Compared with TrkB. FL and control groups, the injection of TrkB. T1 into NAc prolonged the latency of immobility by almost five times, showing an antidepressant-like effect [67].

3.6. LHB

The stimulation of LHB seems can increase the BDNF level in other tissues. A woman diagnosed with MDD treated with deep brain stimulation (DBS) of the LHB, was found an increased level of BDNF in the serum [76]. LHB lesions can improve the inflammatory response of the hippocampus and inhibit the activation of the hippocampal NF- κ B pathway and the expression of inflammatory factors [77]. Inhibiting the function of LHB can increase the expression of BDNF and TrkB in the hippocampus [77, 78].

4. The Downstream and Upstream Molecular of BDNF-TrkB Signaling

The decrease of p-ERK was found in anxiety and depression models [79-81], and MKP-1, the negative regulators of ERK1/2 signaling, was increased [28]. Infusing MEK inhibitor U1026 into the medial prefrontal cortex (mPFC) or dorsal hippocampus (dHP), rats showed anxiety and depression-like behavior, and the expression of p-CREB decreased [82]. As the up-regulation of p-ERK and p-CREB, BDNF increased and the anxiety and depression-like behaviors were alleviated [49, 83].

The decrease of p-CREB was found in anxiety and depression models [84-86], along with hypomyelination and impaired synaptic plasticity [47, 48, 86], and the decreasing BDNF [30, 49]. With the increasing expression of p-CREB after treatment, the myelination was increased, which was confirmed by the increasing MBP, and the synaptic plasticity was improved [47, 86]. The expression of BDNF also increased [87, 88].

The decrease of p-mTOR was found in anxiety and depression models [51, 89, 90], along with decreased synaptic plasticity [90]. After antidepressant treatment, the expression of p-mTOR increased [91, 92], and the therapeutic effect could be inhibited by an mTOR inhibitor [92, 93], blocking the enhancement of excitatory synaptic transmission [32]. Depression-like behavior was observed in mice by unilaterally injecting siRNA into the mPFC to reduce the expression of mTOR. The expression of BDNF in the cortex decreased, and the dysregulation of the release of serotonin and glutamate in the DRN was observed [94].

Increased serum corticosterone, inflammation factors, and oxidative stress status were observed in anxiety and depression models [41, 95, 96], along with the activated astrocyte and microglia [96, 97]. In these models, the

expression of TLR4 and NF- κ B was increased [26, 41, 98]. As the increase of TLR4, TLR4-MyD88-NF- κ B signaling was activated, and the inflammation factors were increased [95, 96]. Anti-inflammation treatment can inhibit the NF- κ B signaling in the LPS-induced model, and alleviate anxiety and depression-like behaviors by decreasing the inflammation factors and the activated astrocyte and microglia and increasing the expression of BDNF [55, 99]. The increase of BDNF in NAc and LHB can lead to depression-like behavior [67, 77] but inhibition of the activated NF- κ B signaling pathway to alleviate the inflammation in NAc and LHB can improve anxiety and depression-like behavior in male mice [77, 100].

5. Conclusions

In this review, we summarized the function of BDNF-TrkB signaling in different brain regions and its antidepressant effect. In all, the inhibition of the BDNF-TrkB signaling pathway in the cortex, hippocampus, hypothalamus, amygdala, and DRN, and the activation of the BDNF-TrkB signaling pathway in ACC, NAc, and LHB is associated with the development of anxiety and depression-like behavior. Studies have shown that the increased BDNF is related to high synaptic plasticity. This indicated that different brain regions have different regulatory effects on anxiety and depression-like behaviors, which requires further exploration. Due to limited studies of BDNF-TrkB signaling in the brain except for the cortex and hippocampus, the role of elevated BDNF-TrkB levels in ACC, NAc, and LHB leading to anxiety and depression-like behaviors remains to be elucidated. The impaired BDNF-TrkB signaling is a mediator, not the cause, of anxiety and depression-like behaviors. It can be regulated by the mTOR, NF- κ B, PGC-1 α -FNDC5, and AMPK signaling pathways. Decreased BDNF makes the body more susceptible to anxiety and depression. BDNF-TrkB signaling pathway plays a key role in antidepressant treatment. Without BDNF, anti-anxiety and anti-depression treatments will not work, and correspondingly, drugs or physical treatment targeting to improve the BDNF-TrkB level are inclined to alleviate the symptom of depression. Therefore, during the treatment of anxiety and depression, efforts can still be made to increase the level of BDNF. But it is still necessary to pay attention to changes in its upstream and downstream pathways in order to accurately treat and prevent the occurrence of anxiety and depression.

Abbreviations

AAV: Adeno-associated virus
 ACC: Anterior cingulate cortex
 Akt: Protein kinase B
 AMPK: AMP-activated protein kinase
 BDNF: Brain-derived neurotrophic factor
 CAT: Catalase
 CeA: Central amygdaloid nucleus
 cKO: Conditional knockout

CORT: Corticosterone
 CREB: cAMP response element binding protein
 CRH: Corticotropin releasing hormone
 CRS: Chronic restraint stress
 CSDS: Chronic social defeat stress
 CUMS: Chronic unpredictable mild stress
 DBS: Deep brain stimulation
 DG: Dentate gyrus
 dHP: Dorsal hippocampus
 DRN: Dorsal raphe nucleus
 EBR: Effort-based reward
 ECT: Electroconvulsive therapy
 EE: Enriched environment
 ERK: Extracellular regulated protein kinases
 EM: Endometriosis model
 FNDC5: Fibronectin type-III domain-containing protein 5
 GFAP: Glial fibrillary acidic protein
 GSH: glutathione
 IBD: Inflammatory bowel disease
 LH: Learned helplessness
 LHb: Lateral habenula
 LPS: Lipopolysaccharide
 MAPK: Mitogen-activated protein kinases
 MBP: Myelin basic protein
 MDA: Malondialdehyde
 MDD: Major depressive disorder
 MEK: Mitogen-activated protein
 MHb: Medial habenula
 mPFC: Medial prefrontal cortex
 mTOR: Mammalian target of rapamycin
 NAc: Nucleus accumbens
 NF- κ B: Nuclear transcription factor- κ B
 Nrf2: NF-E2-related factor 2
 OBX: Olfactory bulbectomy
 PGC-1 α : PPAR- γ coactivator 1 alpha
 PI3K: Phosphatidylinositol-3 kinase
 PSD95: Postsynaptic density protein 95
 PVN: Paraventricular nucleus
 P75^{NTR}: P75 neurotrophin receptor
 SD: Sleep deprivation
 SNI: Spared nerve injury
 SOD: Superoxide dismutase
 SYN: Synaptophysin
 TLR4: Toll-like receptor 4
 tPBM: Transcranial photobiomodulation
 TPO: Thrombopoietin
 VMH: Hypothalamic ventromedial nucleus
 VTA: Ventral tegmental area
 7, 8-DHF: 7, 8-dihydroxyflavone

References

- [1] American Psychiatric Association. (2013). Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition.
- [2] Helen Herrman, Vikram Patel, Christian Kieling, Michael Berk, Claudia Buchweitz, Pim Cuijpers, Toshiaki A Furukawa, Ronald C Kessler, Brandon A Kohrt, Mario Maj, Patrick McGorry, Charles F 3rd Reynolds, Myrna M Weissman, Dixon Chibanda, Christopher Dowrick, Louise M Howard, Christina W Hoven, Martin Knapp, Helen S Mayberg, Miranda Wolpert. (2022). Time for united action on depression: a Lancet-World Psychiatric Association Commission. *Lancet*, 399 (10328), 957-1022. doi: 10.1016/S0140-6736 (21)02141-3.
- [3] John W G Tiller. (2013). Depression and anxiety. *Med J Aust*, 199 (S6), S28-31. doi: 10.5694/mja12.10628.
- [4] G. B. D. Diseases, & Collaborators Injuries. (2020). Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*, 396 (10258), 1204-1222. doi: 10.1016/S0140-6736 (20)30925-9.
- [5] A J Ferrari, A J Somerville, A J Baxter, R Norman, S B Patten, T Vos, & H A Whiteford. (2013). Global variation in the prevalence and incidence of major depressive disorder: a systematic review of the epidemiological literature. *Psychol Med*, 43 (3), 471-481. doi: 10.1017/S0033291712001511.
- [6] Jackson G Thorp, Zachary F Gerring, Lucia Colodro-Conde, Enda M Byrne, Sarah E Medland, Christel M Middeldorp, & Eske M Derks. (2023). The association between trauma exposure, polygenic risk and individual depression symptoms. *Psychiatry Res*, 321, 115101. doi: 10.1016/j.psychres.2023.115101.
- [7] Magdalena Gawrych, Ewelina Cichon, & Andrzej Kiejna. (2022). Depression among young adults - risks and protective factors in the COVID-19 pandemic. *Postep Psychiatr Neurol*, 31 (2), 52-61. doi: 10.5114/ppn.2022.118265.
- [8] Yuanyuan Xiao, Hailiang Ran, Die Fang, Yusan Che, Junwei Peng, Hao Sun, Xuemeng Liang, Lin Chen, Sifan Wang, & Jin Lu. (2022). Childhood maltreatment and depressive disorders in Chinese children and adolescents: A population-based case-control study. *Asian J Psychiatr*, 78, 103312. doi: 10.1016/j.ajp.2022.103312.
- [9] Covid- Mental Disorders Collaborators. (2021). Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet*, 398 (10312), 1700-1712. doi: 10.1016/S0140-6736 (21) 02143-7.
- [10] Gabriel R Fries, Valeria A Saldana, Johannes Finnstein, & Theo Rein. (2023). Molecular pathways of major depressive disorder converge on the synapse. *Mol Psychiatry*, 28 (1), 284-297. doi: 10.1038/s41380-022-01806-1.
- [11] Yiyi Li, Fang Li, Dongdong Qin, Hongyu Chen, Jianhao Wang, Jiabei Wang, Shafei Song, Chao Wang, Yamei Wang, Songyan Liu, Dandan Gao, & Zhi-Hao Wang. (2022). The role of brain derived neurotrophic factor in central nervous system. *Front Aging Neurosci*, 14, 986443. doi: 10.3389/fnagi.2022.986443.
- [12] Daniele Cavaleri, Federico Moretti, Alessandra Bartocetti, Stefano Mauro, Cristina Crocamo, Giuseppe Carra, & Francesco Bartoli. (2023). The role of BDNF in major depressive disorder, related clinical features, and antidepressant treatment: Insight from meta-analyses. *Neurosci Biobehav Rev*, 149, 105159. doi: 10.1016/j.neubiorev.2023.105159.
- [13] Mia Thompson Ray, Cynthia Shannon Weickert, Eugene Wyatt, & Maree J Webster. (2011). Decreased BDNF, trkB-TK+ and GAD67 mRNA expression in the hippocampus of individuals with schizophrenia and mood disorders. *J Psychiatry Neurosci*, 36 (3), 195-203. doi: 10.1503/jpn.100048.

- [14] J-P Guilloux, G Douillard-Guilloux, R Kota, X Wang, A M Gardier, K Martinowich, G C Tseng, D A Lewis, & E. Sibille. (2012). Molecular evidence for BDNF- and GABA-related dysfunctions in the amygdala of female subjects with major depression. *Mol Psychiatry*, 17 (11), 1130-1142. doi: 0.1038/mp.2011.113.
- [15] Giovanni Martinotti, Mauro Pettorruso, Domenico De Berardis, Paola Annunziata Varasano, Gabriella Lucidi Pressanti, Valeria De Remigis, Alessandro Valchera, Valerio Ricci, Marco Di Nicola, Luigi Janiri, Giovanni Biggio, & Massimo Di Giannantonio. (2016). Agomelatine Increases BDNF Serum Levels in Depressed Patients in Correlation with the Improvement of Depressive Symptoms. *Int J Neuropsychopharmacol*, 19 (5). doi: 0.1093/ijnp/pyw003.
- [16] Luisella Bocchio-Chiavetto, Roberta Zanardini, Marco Bortolomasi, Maria Abate, Matilde Segala, Mario Giacomuzzi, Marco Andrea Riva, Eleonora Marchina, Patrizio Pasqualetti, Jorge Perez, & Massimo Gennarelli. (2006). Electroconvulsive Therapy (ECT) increases serum Brain Derived Neurotrophic Factor (BDNF) in drug resistant depressed patients. *Eur Neuropsychopharmacol*, 16 (8), 620-624. doi: 0.1016/j.euroneuro.2006.04.010.
- [17] B Chen, D Dowlathshahi, G M MacQueen, J F Wang, & L T Young. (2001). Increased hippocampal BDNF immunoreactivity in subjects treated with antidepressant medication. *Biol Psychiatry*, 50 (4), 260-265. doi: 0.1016/s0006-3223(01)01083-6.
- [18] Zhiwei Guan, & Jidong Fang. (2006). Peripheral immune activation by lipopolysaccharide decreases neurotrophins in the cortex and hippocampus in rats. *Brain Behav Immun*, 20 (1), 64-71. doi: 0.1016/j.bbi.2005.04.005.
- [19] C R Yang, Y Y Bai, C S Ruan, F H Zhou, F Li, C Q Li, & X F Zhou. (2017). Injection of Anti-proBDNF in Anterior Cingulate Cortex (ACC) Reverses Chronic Stress-Induced Adverse Mood Behaviors in Mice. *Neurotox Res*, 31 (2), 298-308. doi: 0.1007/s12640-016-9687-4.
- [20] M Nibuya, S Morinobu, & R S Duman. (1995). Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments. *J Neurosci*, 15 (11), 7539-7547. doi: 0.1523/JNEUROSCI.15-11-07539.1995.
- [21] Yafei Ji, Jie Luo, Jiuseng Zeng, Yang Fang, Rong Liu, Fei Luan, & Nan Zeng. (2021). Xiaoyao Pills Ameliorate Depression-like Behaviors and Oxidative Stress Induced by Olfactory Bulbectomy in Rats via the Activation of the PIK3CA-AKT1-NFE2L2/BDNF Signaling Pathway. *Front Pharmacol*, 12, 643456. doi: 0.3389/fphar.2021.643456.
- [22] Indira Mendez-David, Laurent Tritschler, Zeina El Ali, Marie-Helene Damiens, Marc Pallardy, Denis J David, Saadia Kerdine-Romer, & Alain M Gardier. (2015). Nrf2-signaling and BDNF: A new target for the antidepressant-like activity of chronic fluoxetine treatment in a mouse model of anxiety/depression. *Neurosci Lett*, 597, 121-126. doi: 0.1016/j.neulet.2015.04.036.
- [23] Chun Yang, Yukihiko Shirayama, Ji-Chun Zhang, Qian Ren, & Kenji Hashimoto. (2015). Regional differences in brain-derived neurotrophic factor levels and dendritic spine density confer resilience to inescapable stress. *Int J Neuropsychopharmacol*, 18 (7), pyu121. doi: 0.1093/ijnp/pyu121.
- [24] Lucia Recinella, Annalisa Chiavaroli, Giustino Orlando, Claudio Ferrante, Serena Veschi, Alessandro Cama, Guya Diletta Marconi, Francesca Diomede, Iacopo Gesmundo, Riccarda Granata, Renzhi Cai, Wei Sha, Andrew V Schally, Luigi Brunetti, & Sheila Leone. (2021). Effects of growth hormone-releasing hormone receptor antagonist MIA-602 in mice with emotional disorders: a potential treatment for PTSD. *Mol Psychiatry*, 26 (12), 7465-7474. doi: 0.1038/s41380-021-01228-5.
- [25] Ji-chun Zhang, Jin Wu, Yuko Fujita, Wei Yao, Qian Ren, Chun Yang, Su-xia Li, Yukihiko Shirayama, & Kenji Hashimoto. (2014). Antidepressant effects of TrkB ligands on depression-like behavior and dendritic changes in mice after inflammation. *Int J Neuropsychopharmacol*, 18 (4). doi: 0.1093/ijnp/pyu077.
- [26] Parvaneh Mohseni-Moghaddam, Manijeh Dogani, Motahare Hatami, Samira Roohollahi, Azam Amiresmaeli, & Nayereh Askari. (2022). A behavioral and molecular study; ameliorated anxiety-like behavior and cognitive dysfunction in a rat model of chronic unpredictable stress treated with oregano extract. *Brain Behav*, 12 (8), e2727. doi: 0.1002/brb3.2727.
- [27] Sinead M Gibney, Barry McGuinness, Christine Prendergast, Andrew Harkin, & Thomas J Connor. (2013). Poly I: C-induced activation of the immune response is accompanied by depression and anxiety-like behaviours, kynurenine pathway activation and reduced BDNF expression. *Brain Behav Immun*, 28, 170-181. doi: 0.1016/j.bbi.2012.11.010.
- [28] Qin Ru, Qi Xiong, Mei Zhou, Lin Chen, Xiang Tian, Huqiao Xiao, Chaoying Li, & Ying Li. (2019). Withdrawal from chronic treatment with methamphetamine induces anxiety and depression-like behavior in mice. *Psychiatry Res*, 271, 476-483. doi: 0.1016/j.psychres.2018.11.072.
- [29] Adnan Khan, Bushra Shal, Muhammad Naveed, Bakht Nasir, Nadeem Irshad, Hussain Ali, & Salman Khan. (2020). Matrine alleviates neurobehavioral alterations via modulation of JNK-mediated caspase-3 and BDNF/VEGF signaling in a mouse model of burn injury. *Psychopharmacology (Berl)*, 237 (8), 2327-2343. doi: 0.1007/s00213-020-05537-5.
- [30] Basma Youssef, Kholoud S Ramadan, Shaimaa ElShebiney, & Ehab A Ibrahim. (2022). Antidepressant-like effects of aqueous extracts of miswak (*Salvadora persica*) and date palm (*Phoenix dactylifera*) on depression-like behaviors using CUMS model in male rats. *J Food Biochem*, 46 (8), e14164. doi: 0.1111/jfbc.14164.
- [31] Lanwei Hou, Yirui Qi, Hongwei Sun, Gang Wang, Qi Li, Yanyu Wang, Zuoji Zhang, Zhongde Du, & Lin Sun. (2018). Applying ketamine to alleviate the PTSD-like effects by regulating the HCN1-related BDNF. *Prog Neuropsychopharmacol Biol Psychiatry*, 86, 313-321. doi: 0.1016/j.pnpbp.2018.03.019.
- [32] Fei Guo, Bing Zhang, Zhiwen Fu, Yuqin Ma, Yu Gao, Fuyi Shen, Chenggang Huang, & Yang Li. (2016). The rapid antidepressant and anxiolytic-like effects of YY-21 involve enhancement of excitatory synaptic transmission via activation of mTOR signaling in the mPFC. *Eur Neuropsychopharmacol*, 26 (7), 1087-1098. doi: 0.1016/j.euroneuro.2016.05.006.
- [33] Chia-Yu Chang, How-Ran Guo, Wan-Chen Tsai, Kai-Lin Yang, Li-Chuan Lin, Tain-Junn Cheng, & Jiunn-Jye Chuu. (2015). Subchronic Arsenic Exposure Induces Anxiety-Like Behaviors in Normal Mice and Enhances Depression-Like Behaviors in the Chemically Induced Mouse Model of Depression. *Biomed Res Int*, 2015, 159015. doi: 0.1155/2015/159015.

- [34] Raffaella Alo, Maria Mele, Gilda Fazzari, Ennio Avolio, & Marcello Canonaco. (2015). Exposure to sub-chronic unpredictable stress accounts for antidepressant-like effects in hamsters treated with BDNF and CNQX. *Brain Res Bull*, 118, 65-77. doi: 0.1016/j.brainresbull.2015.09.007.
- [35] P A Lapchak, D M Araujo, & F. Hefti. (1993). Systemic interleukin-1 beta decreases brain-derived neurotrophic factor messenger RNA expression in the rat hippocampal formation. *Neuroscience*, 53 (2), 297-301. doi: 0.1016/0306-4522(93)90196-m.
- [36] Ying Yu, Zhaokun Shi, Dan Xu, Ying Li, Jun Qin, Zhaohui Zhang, & Hui Wang. (2020). Prenatal ethanol exposure increases susceptibility to depression- and anxiety-like behavior in adult female offspring and its underlying mechanism. *Reprod Toxicol*, 96, 36-46. doi: 0.1016/j.reprotox.2020.05.015.
- [37] Ga-Young Choi, Hyun-Bum Kim, Eun-Sang Hwang, Seok Lee, Min-Ji Kim, Ji-Young Choi, Sung-Ok Lee, Sang-Seong Kim, & Ji-Ho Park. (2017). Curcumin Alters Neural Plasticity and Viability of Intact Hippocampal Circuits and Attenuates Behavioral Despair and COX-2 Expression in Chronically Stressed Rats. *Mediators Inflamm*, 2017, 6280925. doi: 0.1155/2017/6280925.
- [38] Paulo Wagner Linhares Lima Filho, Adriano Jose Maia Chaves Filho, Charliene Freire Xavier Vieira, Tatiana de Queiroz Oliveira, Michelle Verde Ramo Soares, Paloma Marinho Juca, Joao Quevedo, Tatiana Barichello, Danielle Macedo, & Francisco das Chagas Medeiros. (2019). Peritoneal endometriosis induces time-related depressive- and anxiety-like alterations in female rats: involvement of hippocampal pro-oxidative and BDNF alterations. *Metab Brain Dis*, 34 (3), 909-925. doi: 0.1007/s11011-019-00397-1.
- [39] Haidar Tafner Curi, Clarissa Tavares Dias, Maria Laura Monteiro da Luz Camargo, Paula Dos Santos Gomez, Moises Felipe Pereira Gomes, Jose Ivo Araujo Beserra-Filho, Alessandra Medeiros, Alessandra Mussi Ribeiro, Fernando Moreira Simabuco, Rafael Herling Lambertucci, & Cristiano Mendes-da-Silva. (2021). Maternal high-fat diet increases anhedonic behavior and modulates hippocampal Mash1 and BDNF expression in adult offspring. *Neurosci Lett*, 764, 136239. doi: 0.1016/j.neulet.2021.136239.
- [40] Lorrane Kelle da Silva Moreira, Adriane Ferreira de Brito, Dayane Moreira da Silva, Lorryne Siqueira, Daiany Priscilla Bueno da Silva, Carina Sofia Cardoso, Iziara Ferreira Florentino, Pablinny Moreira Galdino de Carvalho, Paulo Cesar Ghedini, Ricardo Menegatti, & Elson Alves Costa. (2021). Potential antidepressant-like effect of piperazine derivative LQFM212 in mice: Role of monoaminergic pathway and brain-derived neurotrophic factor. *Behav Brain Res*, 401, 113066. doi: 0.1016/j.bbr.2020.113066.
- [41] Arya Haj-Mirzaian, Shayan Amiri, Hossein Amini-Khoei, Mir-Jamal Hosseini, Arvin Haj-Mirzaian, Majid Momeny, Maryam Rahimi-Balaei, & Ahmad Reza Dehpour. (2017). Anxiety- and Depressive-Like Behaviors are Associated with Altered Hippocampal Energy and Inflammatory Status in a Mouse Model of Crohn's Disease. *Neuroscience*, 366, 124-137. doi: 0.1016/j.neuroscience.2017.10.023.
- [42] Xiaoyu Song, Bo Liu, Lingyu Cui, Biao Zhou, Weiwei Liu, Fanxing Xu, Toshihiko Hayashi, Shunji Hattori, Yuko Ushiki-Kaku, Shin-Ichi Tashiro, & Takashi Ikejima. (2017). Silibinin ameliorates anxiety/depression-like behaviors in amyloid beta-treated rats by upregulating BDNF/TrkB pathway and attenuating autophagy in hippocampus. *Physiol Behav*, 179, 487-493. doi: 0.1016/j.physbeh.2017.07.023.
- [43] Jie Liu, Chen Yang, Jing Yang, Xiaojie Song, Wei Han, Mingdan Xie, Li Cheng, Lingling Xie, Hengsheng Chen, & Li Jiang. (2019). Effects of early postnatal exposure to fine particulate matter on emotional and cognitive development and structural synaptic plasticity in immature and mature rats. *Brain Behav*, 9 (12), e01453. doi: 0.1002/brb3.1453.
- [44] Young-Ju Lee, Hye Ryeong Kim, Chang Youn Lee, Sung-Ae Hyun, Moon Yi Ko, Byoung-Seok Lee, Dae Youn Hwang, & Minhan Ka. (2020). 2-Phenylethylamine (PEA) Ameliorates Corticosterone-Induced Depression-Like Phenotype via the BDNF/TrkB/CREB Signaling Pathway. *Int J Mol Sci*, 21 (23). doi: 0.3390/ijms21239103.
- [45] Narmin Farazi, Javad Mahmoudi, Saeed Sadigh-Eteghad, Fereshteh Farajdokht, & Seyed Hossein Rasta. (2022). Synergistic effects of combined therapy with transcranial photobiomodulation and enriched environment on depressive- and anxiety-like behaviors in a mice model of noise stress. *Lasers Med Sci*, 37 (2), 1181-1191. doi: 0.1007/s10103-021-03370-6.
- [46] Pit Shan Chong, Chi Him Poon, Jaydeep Roy, Ka Chun Tsui, Sze Yuen Lew, Michael Weng Lok Phang, Rachael Julia Yuenyinn Tan, Poh Guat Cheng, Man-Lung Fung, Kah Hui Wong, & Lee Wei Lim. (2021). Neurogenesis-dependent antidepressant-like activity of *Hericium erinaceus* in an animal model of depression. *Chin Med*, 16 (1), 132. doi: 0.1186/s13020-021-00546-8.
- [47] Wen-Xue Liu, Jing Wang, Ze-Min Xie, Ning Xu, Guang-Fen Zhang, Min Jia, Zhi-Qiang Zhou, Kenji Hashimoto, & Jian-Jun Yang. (2016). Regulation of glutamate transporter 1 via BDNF-TrkB signaling plays a role in the anti-apoptotic and antidepressant effects of ketamine in chronic unpredictable stress model of depression. *Psychopharmacology (Berl)*, 233 (3), 405-415. doi: 0.1007/s00213-015-4128-2.
- [48] Bo He, Dan Xu, Chong Zhang, Li Zhang, & Hui Wang. (2018). Prenatal food restriction induces neurobehavioral abnormalities in adult female offspring rats and alters intrauterine programming. *Toxicol Res (Camb)*, 7 (2), 293-306. doi: 0.1039/c7tx00133a.
- [49] Tingxu Yan, Yingying Sun, Feng Xiao, Bo Wu, Kaishun Bi, Bosai He, & Ying Jia. (2019). *Schisandrae Chinensis Fructus* inhibits behavioral deficits induced by sleep deprivation and chronic unpredictable mild stress via increased signaling of brain-derived neurotrophic factor. *Phytother Res*, 33 (12), 3177-3190. doi: 0.1002/ptr.6489.
- [50] Ayyub Babaei, Maryam Nourshahi, Maryam Fani, Zahra Entezari, Seyed Behnamedin Jameie, & Abbas Haghparsat. (2021). The effectiveness of continuous and interval exercise preconditioning against chronic unpredictable stress: Involvement of hippocampal PGC-1alpha/FNDC5/BDNF pathway. *J Psychiatr Res*, 136, 173-183. doi: 0.1016/j.jpsychires.2021.02.006.
- [51] Julia M Rosa, Francis L Pazini, Gislaire Olescowicz, Anderson Camargo, Morgana Moretti, Joana Gil-Mohapel, & Ana Lucia S Rodrigues. (2019). Prophylactic effect of physical exercise on Abeta (1-40)-induced depressive-like behavior: Role of BDNF, mTOR signaling, cell proliferation and survival in the hippocampus. *Prog Neuropsychopharmacol Biol Psychiatry*, 94, 109646. doi: 0.1016/j.pnpbp.2019.109646.

- [52] Osamu Nakagawasai, Kotaro Yamada, Kohei Takahashi, Takayo Odaira, Wakana Sakuma, Daisuke Ishizawa, Naruya Takahashi, Kentaro Onuma, Chikako Hozumi, Wataru Nemoto, & Koichi Tan-No. (2020). Liver hydrolysate prevents depressive-like behavior in an animal model of colitis: Involvement of hippocampal neurogenesis via the AMPK/BDNF pathway. *Behav Brain Res*, 390, 112640. doi: 0.1016/j.bbr.2020.112640.
- [53] Payal Bajaj, & Gurcharan Kaur. (2022). Acute Sleep Deprivation-Induced Anxiety and Disruption of Hypothalamic Cell Survival and Plasticity: A Mechanistic Study of Protection by Butanol Extract of *Tinospora cordifolia*. *Neurochem Res*, 47 (6), 1692-1706. doi: 0.1007/s11064-022-03562-8.
- [54] Jing Wen, Yaowei Xu, Zhixiang Yu, Yifan Zhou, Wenting Wang, Jingjie Yang, Yiming Wang, Qian Bai, & Zhisong Li. (2022). The cAMP Response Element-Binding Protein/Brain-Derived Neurotrophic Factor Pathway in Anterior Cingulate Cortex Regulates Neuropathic Pain and Anxiodepression Like Behaviors in Rats. *Front Mol Neurosci*, 15, 831151. doi: 0.3389/fnmol.2022.831151.
- [55] Abimbola S Olugbemide, Benneth Ben-Azu, Adewale G Bakre, Abayomi M Ajayi, Omowumi Femi-Akinlosotu, & Solomon Umukoro. (2021). Naringenin improves depressive- and anxiety-like behaviors in mice exposed to repeated hypoxic stress through modulation of oxido-inflammatory mediators and NF-kB/BDNF expressions. *Brain Res Bull*, 169, 214-227. doi: 0.1016/j.brainresbull.2020.12.003.
- [56] Su Yeon Seo, Se Kyun Bang, Suk Yun Kang, Seong Jin Cho, Kwang Ho Choi, & Yeon Hee Ryu. (2021). Acupuncture Alleviates Anxiety and 22-kHz Ultrasonic Vocalizations in Rats Subjected to Repeated Alcohol Administration by Modulating the Brain-Derived Neurotrophic Factor/Corticotropin-Releasing Hormone Signaling Pathway. *Int J Mol Sci*, 22 (8). doi: 0.3390/ijms22084037.
- [57] Olivier Berton, Colleen A McClung, Ralph J Dileone, Vaishnav Krishnan, William Renthal, Scott J Russo, Danielle Graham, Nadia M Tsankova, Carlos A Bolanos, Maribel Rios, Lisa M Monteggia, David W Self, & Eric J Nestler. (2006). Essential role of BDNF in the mesolimbic dopamine pathway in social defeat stress. *Science*, 311 (5762), 864-868. doi: 0.1126/science.1120972.
- [58] J A Siuciak, D R Lewis, S J Wiegand, & R M Lindsay. (1997). Antidepressant-like effect of brain-derived neurotrophic factor (BDNF). *Pharmacol Biochem Behav*, 56 (1), 131-137. doi: 0.1016/S0091-3057 (96)00169-4.
- [59] Ting Lei, Dan Dong, Meiying Song, Yanfei Sun, Xiaofeng Liu, & Hua Zhao. (2020). Risperidone treatment in the lateral habenula improves despair-like behavior in mice. *Neuropsychopharmacology*, 45 (10), 1717-1724. doi: 0.1038/s41386-020-0652-9.
- [60] Mannan Abdul, Hao-Qi Yan, Wei-Nan Zhao, Xiao-Bin Lyu, Zheng Xu, Xiao-Lu Yu, Yi-Hong Gao, & Jun-Li Cao. (2022). VTA-NAc glutaminergic projection involves in the regulation of pain and pain-related anxiety. *Front Mol Neurosci*, 15, 1083671. doi: 0.3389/fnmol.2022.1083671.
- [61] Daniel C Lowes, Linda A Chamberlin, Lisa N Kretsge, Emma S Holt, Atheer I Abbas, Alan J Park, Lyubov Yusufova, Zachary H Bretton, Ayesha Firdous, Armen G Enikolopov, Joshua A Gordon, & Alexander Z Harris. (2021). Ventral tegmental area GABA neurons mediate stress-induced blunted reward-seeking in mice. *Nat Commun*, 12 (1), 3539. doi: 0.1038/s41467-021-23906-2.
- [62] Megumi Adachi, Anita E Autry, Melissa Mahgoub, Kanzo Suzuki, & Lisa M Monteggia. (2017). TrkB Signaling in Dorsal Raphe Nucleus is Essential for Antidepressant Efficacy and Normal Aggression Behavior. *Neuropsychopharmacology*, 42 (4), 886-894. doi: 0.1038/npp.2016.201.
- [63] Emily Ploppert, Joanna Jacob, Ana Deutsch, Sally Watanabe, Katherine Gillenwater, Alison Choe, George B Cruz, Ericka Cabanas, Michelle A Vasquez, Zaid Ayaz, Lorenz S Neuwirth, & Kelly Lambert. (2022). Influence of Effort-based Reward Training on Neuroadaptive Cognitive Responses: Implications for Preclinical Behavioral Approaches for Depressive Symptoms. *Neuroscience*, 500, 63-78. doi: 0.1016/j.neuroscience.2022.08.002.
- [64] M Kent, S Scott, S Lambert, E Kirk, B Terhune-Cotter, B Thompson, S Neal, B Dozier, M Bardi, & K. Lambert. (2018). Contingency Training Alters Neurobiological Components of Emotional Resilience in Male and Female Rats. *Neuroscience*, 386, 121-136. doi: 0.1016/j.neuroscience.2018.06.010.
- [65] Megumi Adachi, Michel Barrot, Anita E Autry, David Theobald, & Lisa M Monteggia. (2008). Selective loss of brain-derived neurotrophic factor in the dentate gyrus attenuates antidepressant efficacy. *Biol Psychiatry*, 63 (7), 642-649. doi: 0.1016/j.biopsych.2007.09.019.
- [66] Yukihiko Shirayama, Andrew C-H Chen, Shin Nakagawa, David S Russell, & Ronald S Duman. (2002). Brain-derived neurotrophic factor produces antidepressant effects in behavioral models of depression. *J Neurosci*, 22 (8), 3251-3261. doi: 0.1523/JNEUROSCI.22-08-03251.2002.
- [67] Amelia J Eisch, Carlos A Bolanos, Joris de Wit, Ryan D Simonak, Cindy M Pudiak, Michel Barrot, Joost Verhaagen, & Eric J Nestler. (2003). Brain-derived neurotrophic factor in the ventral midbrain-nucleus accumbens pathway: a role in depression. *Biol Psychiatry*, 54 (10), 994-1005. doi: 0.1016/j.biopsych.2003.08.003.
- [68] K R Jones, I Farinas, C Backus, & L F Reichardt. (1994). Targeted disruption of the BDNF gene perturbs brain and sensory neuron development but not motor neuron development. *Cell*, 76 (6), 989-999. doi: 0.1016/0092-8674 (94)90377-8.
- [69] Tommi Saarelainen, Panu Hendolin, Guilherme Lucas, Eija Koponen, Mikko Sairanen, Ewen MacDonald, Karin Agerman, Annakaisa Haapasalo, Hiroyuki Nawa, Raquel Aloyz, Patrik Ernfors, & Eero Castren. (2003). Activation of the TrkB neurotrophin receptor is induced by antidepressant drugs and is required for antidepressant-induced behavioral effects. *J Neurosci*, 23 (1), 349-357. doi: 0.1523/JNEUROSCI.23-01-00349.2003.
- [70] Lisa M Monteggia, Bryan Luikart, Michel Barrot, David Theobald, Irena Malkovska, Serge Nef, Luis F Parada, & Eric J Nestler. (2007). Brain-derived neurotrophic factor conditional knockouts show gender differences in depression-related behaviors. *Biol Psychiatry*, 61 (2), 187-197. doi: 0.1016/j.biopsych.2006.03.021.
- [71] Lisa M Monteggia, Michel Barrot, Craig M Powell, Olivier Berton, Victor Galanis, Terry Gemelli, Sven Meuth, Andreas Nagy, Robert W Greene, & Eric J Nestler. (2004). Essential role of brain-derived neurotrophic factor in adult hippocampal function. *Proc Natl Acad Sci U S A*, 101 (29), 10827-10832. doi: 0.1073/pnas.0402141101.
- [72] Keri Martinowich, Hussein Manji, & Bai Lu. (2007). New insights into BDNF function in depression and anxiety. *Nat Neurosci*, 10 (9), 1089-1093. doi: 0.1038/nn1971.

- [73] T Saarelainen, J A Lukkarinen, S Koponen, O H Grohn, J Jolkonen, E Koponen, A Haapasalo, L Alhonen, G Wong, J Koistinaho, R A Kauppinen, & E. Castren. (2000). Transgenic mice overexpressing truncated trkB neurotrophin receptors in neurons show increased susceptibility to cortical injury after focal cerebral ischemia. *Mol Cell Neurosci*, 16 (2), 87-96. doi: 0.1006/mcne.2000.0863.
- [74] Xiao Zhuang, Bing Zhan, Yufeng Jia, Chaoze Li, Nan Wu, Ming Zhao, Nuo Chen, Yaxin Guo, Yingxin Du, Yi Zhang, Baihui Cao, Yan Li, Faliang Zhu, Chun Guo, Qun Wang, Yuan Li, & Lining Zhang. (2022). IL-33 in the basolateral amygdala integrates neuroinflammation into anxiogenic circuits via modulating BDNF expression. *Brain Behav Immun*, 102, 98-109. doi: 0.1016/j.bbi.2022.02.019.
- [75] O Guillin, J Diaz, P Carroll, N Griffon, J C Schwartz, & P. Sokoloff. (2001). BDNF controls dopamine D3 receptor expression and triggers behavioural sensitization. *Nature*, 411 (6833), 86-89. doi: 0.1038/35075076.
- [76] Carolin Hoyer, Laura Kranaster, Alexander Sartorius, Rainer Hellweg, & Peter Gass. (2012). Long-term course of brain-derived neurotrophic factor serum levels in a patient treated with deep brain stimulation of the lateral habenula. *Neuropsychobiology*, 65 (3), 147-152. doi: 0.1159/000335243.
- [77] Yachun Wang, Peng Qu, Yimeng Sun, Ziang Li, Lei Liu, & Limin Yang. (2022). Association between increased inflammatory cytokine expression in the lateral habenular nucleus and depressive-like behavior induced by unpredictable chronic stress in rats. *Exp Neurol*, 349, 113964. doi: 0.1016/j.expneurol.2021.113964.
- [78] Peng Qu, Yachun Wang, Lei Liu, Mengmeng Qi, Yimeng Sun, Siyang Zheng, Zichen Xu, Changhong Liu, Xiaoyan Bai, Qinggao Zhang, & Limin Yang. (2020). Habenula lesions improve glucose metabolism in rats with type 2 diabetes by increasing insulin sensitivity and inhibiting gluconeogenesis. *BMJ Open Diabetes Res Care*, 8 (1). doi: 0.1136/bmjdr-2020-001250.
- [79] Luciano Roman-Albasini, Gabriela Diaz-Veliz, Felipe Antonio Olave, Felipe Ignacio Aguayo, Gonzalo Garcia-Rojas, Wladimir Antonio Corrales, Juan Pablo Silva, Ana Maria Avalos, Paulina S Rojas, Esteban Aliaga, & Jenny Lucy Fiedler. (2020). Antidepressant-relevant behavioral and synaptic molecular effects of long-term fasudil treatment in chronically stressed male rats. *Neurobiol Stress*, 13, 100234. doi: 0.1016/j.ynstr.2020.100234.
- [80] Ya Li, Yajing Chen, Xiaoxiao Gao, & Zhongqiu Zhang. (2017). The behavioral deficits and cognitive impairment are correlated with decreased IGF-II and ERK in depressed mice induced by chronic unpredictable stress. *Int J Neurosci*, 127 (12), 1096-1103. doi: 0.1080/00207454.2017.1337014.
- [81] Yvfeng Liang, Jiahong Li, Tao Jin, Ting Gu, Qingjie Zhu, Yizhong Hu, Yang Yang, Jisui Li, Donghong Wu, Kesheng Jiang, & Xiaohong Xu. (2018). Bisphenol-A inhibits improvement of testosterone in anxiety- and depression-like behaviors in gonadectomized male mice. *Horm Behav*, 102, 129-138. doi: 0.1016/j.yhbeh.2018.05.012.
- [82] Xiaoli Qi, Wenjuan Lin, Donglin Wang, Yuqin Pan, Weiwen Wang, & Meng Sun. (2009). A role for the extracellular signal-regulated kinase signal pathway in depressive-like behavior. *Behav Brain Res*, 199 (2), 203-209. doi: 0.1016/j.bbr.2008.11.051.
- [83] Vittoria Borgonetti, Francisco Les, Victor Lopez, & Nicoletta Galeotti. (2020). Attenuation of Anxiety-Like Behavior by *Helichrysum stoechas* (L.) Moench Methanolic Extract through Up-Regulation of ERK Signaling Pathways in Noradrenergic Neurons. *Pharmaceuticals (Basel)*, 13 (12). doi: 0.3390/ph13120472.
- [84] Zhen Wu, Pengyu Wang, Daodong Pan, Xiaoqun Zeng, Yuxing Guo, & Guangsheng Zhao. (2021). Effect of adzuki bean sprout fermented milk enriched in gamma-aminobutyric acid on mild depression in a mouse model. *J Dairy Sci*, 104 (1), 78-91. doi: 0.3168/jds.2020-19154.
- [85] Afzal Misrani, Sidra Tabassum, Xi Chen, Shu-Yi Tan, Ji-Chen Wang, Li Yang, & Cheng Long. (2019). Differential effects of citalopram on sleep-deprivation-induced depressive-like behavior and memory impairments in mice. *Prog Neuropsychopharmacol Biol Psychiatry*, 88, 102-111. doi: 0.1016/j.pnpbp.2018.07.013.
- [86] Gangadharan Thamizhoviya, & Arambakkam Janardhanam Vanisree. (2019). Enriched environment modulates behavior, myelination and augments molecules governing the plasticity in the forebrain region of rats exposed to chronic immobilization stress. *Metab Brain Dis*, 34 (3), 875-887. doi: 0.1007/s11011-018-0370-8.
- [87] Xian-Feng Huang, Wen-Tao Jiang, Li Liu, Fang-Chen Song, Xia Zhu, Gui-Lan Shi, Shu-Ming Ding, Heng-Ming Ke, Wei Wang, James M O'Donnell, Han-Ting Zhang, Hai-Bin Luo, Yi-Qian Wan, Guo-Qiang Song, & Ying Xu. (2018). A novel PDE9 inhibitor WYQ-C36D ameliorates corticosterone-induced neurotoxicity and depression-like behaviors by cGMP-CREB-related signaling. *CNS Neurosci Ther*, 24 (10), 889-896. doi: 0.1111/cns.12864.
- [88] Xi-Dan Zhou, Yu Zheng, Rakesh Sharma, & Zhang-Jin Zhang. (2019). Total Polysaccharides of Lily Bulb Ameliorate Menopause-Like Behavior in Ovariectomized Mice: Multiple Mechanisms Distinct from Estrogen Therapy. *Oxid Med Cell Longev*, 2019, 6869350. doi: 0.1155/2019/6869350.
- [89] Arindam Ghosh Mazumder, Pallavi Sharma, Vikram Patial, & Damanpreet Singh. (2017). Ginkgo biloba L. attenuates spontaneous recurrent seizures and associated neurological conditions in lithium-pilocarpine rat model of temporal lobe epilepsy through inhibition of mammalian target of rapamycin pathway hyperactivation. *J Ethnopharmacol*, 204, 8-17. doi: 0.1016/j.jep.2017.03.060.
- [90] Anfeng Wang, Xiaojuan Zou, Jiajia Wu, Qingyu Ma, Naijun Yuan, Fengmin Ding, Xiaojuan Li, & Jiaxu Chen. (2020). Early-Life Stress Alters Synaptic Plasticity and mTOR Signaling: Correlation With Anxiety-Like and Cognition-Related Behavior. *Front Genet*, 11, 590068. doi: 0.3389/fgene.2020.590068.
- [91] Xiuping Sun, Xianglei Li, Ruile Pan, Yanfeng Xu, Qiong Wang, & Mingjing Song. (2018). Total Saikosaponins of *Bupleurum yinchowense* reduces depressive, anxiety-like behavior and increases synaptic proteins expression in chronic corticosterone-treated mice. *BMC Complement Altern Med*, 18 (1), 117. doi: 0.1186/s12906-018-2186-9.
- [92] Ju-Young Oh, Yu-Kang Kim, Seung-Nam Kim, Bombi Lee, Jae-Hwan Jang, Sunoh Kwon, & Hi-Joon Park. (2018). Acupuncture modulates stress response by the mTOR signaling pathway in a rat post-traumatic stress disorder model. *Sci Rep*, 8 (1), 11864. doi: 0.1038/s41598-018-30337-5.

- [93] Ju-Young Oh, Quan Feng Liu, Cai Hua, Ha Jin Jeong, Jae-Hwan Jang, Songhee Jeon, & Hi-Joon Park. (2020). Intranasal Administration of Melanin-Concentrating Hormone Reduces Stress-Induced Anxiety- and Depressive-Like Behaviors in Rodents. *Exp Neurol*, 29 (6), 453-469. doi: 0.5607/en20024.
- [94] Emilio Garro-Martinez, Maria Neus Fullana, Eva Florensa-Zanuy, Julia Senserrich, Veronica Paz, Esther Ruiz-Bronchal, Albert Adell, Elena Castro, Alvaro Diaz, Angel Pazos, Analia Bortolozzi, & Fuencisla Pilar-Cuellar. (2021). mTOR Knockdown in the Infralimbic Cortex Evokes A Depressive-like State in Mouse. *Int J Mol Sci*, 22 (16). doi: 0.3390/ijms22168671.
- [95] Feini Zhou, Hao Jiang, Ning Kong, Jiangnan Lin, Fan Zhang, Ting Mai, Zhijian Cao, & Maosheng Xu. (2022). Electroacupuncture Attenuated Anxiety and Depression-Like Behavior via Inhibition of Hippocampal Inflammatory Response and Metabolic Disorders in TNBS-Induced IBD Rats. *Oxid Med Cell Longev*, 2022, 8295580. doi: 0.1155/2022/8295580.
- [96] Walaa Yehia Abdelzاهر, Hanaa H Mohammed, Nermeen N Welson, Gaber El-Saber Batiha, Roua S Baty, & Asmaa Mohamed Abdel-Aziz. (2021). Rivaroxaban Modulates TLR4/Myd88/NF-Kbeta Signaling Pathway in a Dose-Dependent Manner With Suppression of Oxidative Stress and Inflammation in an Experimental Model of Depression. *Front Pharmacol*, 12, 715354. doi: 0.3389/fphar.2021.715354.
- [97] Xi Xiao, Hui Zhang, Wen Ning, Zhuo Yang, Yue Wang, & Tao Zhang. (2022). Knockdown of FSTL1 inhibits microglia activation and alleviates depressive-like symptoms through modulating TLR4/MyD88/NF-kappaB pathway in CUMS mice. *Exp Neurol*, 353, 114060. doi: 0.1016/j.expneurol.2022.114060.
- [98] Meng Wang, Li-Rong Feng, Zi-Long Li, Kai-Ge Ma, Ke-Wei Chang, Xin-Lin Chen, Peng-Bo Yang, Sheng-Feng Ji, Yan-Bing Ma, Hua Han, John Bosco Ruganzua, Wei-Na Yang, & Yi-Hua Qian. (2021). Thymosin beta4 reverses phenotypic polarization of glial cells and cognitive impairment via negative regulation of NF-kappaB signaling axis in APP/PS1 mice. *J Neuroinflammation*, 18 (1), 146. doi: 0.1186/s12974-021-02166-3.
- [99] Fernanda Severo Sabedra Sousa, Paloma Taborda Birmann, Suely Ribeiro Bampi, Mariana G Fronza, Renata Balaguez, Diego Alves, Marlon Regis Leite, Cristina Wayne Nogueira, Cesar Augusto Bruning, & Lucielli Savegnago. (2019). Lipopolysaccharide-induced depressive-like, anxiogenic-like and hyperalgesic behavior is attenuated by acute administration of alpha-(phenylselanyl) acetophenone in mice. *Neuropharmacology*, 146, 128-137. doi: 0.1016/j.neuropharm.2018.11.028.
- [100] Lea Decarie-Spain, Sandeep Sharma, Cecile Hryhorczuk, Victor Issa-Garcia, Philip A Barker, Nathalie Arbour, Thierry Alquier, & Stephanie Fulton. (2018). Nucleus accumbens inflammation mediates anxiodepressive behavior and compulsive sucrose seeking elicited by saturated dietary fat. *Mol Metab*, 10, 1-13. doi: 0.1016/j.molmet.2018.01.018.